

What are the radio outflows in gamma-ray binaries?

Valentí Bosch-Ramon

Universitat de Barcelona/Max Planck Institut für Kernphysik

High Energy Phenomena in Relativistic Outflows II
Buenos Aires, Argentina

27/10/2009

Outline

- 1 Introduction
- 2 Exploring the scenario
- 3 Flux and spectral evolution in LS 5039
- 4 Morphological evolution
- 5 Summary

Outline

- 1 Introduction
- 2 Exploring the scenario
- 3 Flux and spectral evolution in LS 5039
- 4 Morphological evolution
- 5 Summary

Gamma-ray binaries

- PSR B1258–63: Be + young non-accreting pulsar + radio (Aharonian et al. 2005)
- LS 5039: O ms + pulsar or black hole + radio (Aharonian et al. 2005)
- LS I +61 303: Be + pulsar or black hole + radio (Albert et al. 2006; Acciari et al. 2008)
- Cygnus X-1: O Sg + black hole + radio (Albert et al. 2007)
- HESS J0632+057?: massive star? + pulsar or black hole? + radio (Hinton et al. 2009)

Is radio emission produced by a relativistic outflow?


Pair creation and radio emission

- Gamma-rays ≥ 100 GeV are absorbed by the stellar photons producing e^\pm pairs.
- Pairs are quickly trapped in the stellar wind magnetic field.
- Radio and X-ray emission from the pairs in the wind may be relevant (Bosch-Ramon, Khangulyan & Aharonian 2008).
- Flux, lightcurve, and morphology in the radio band will be linked to the gamma-ray emitter and the stellar wind properties.

Outline

- 1 Introduction
- 2 Exploring the scenario
- 3 Flux and spectral evolution in LS 5039
- 4 Morphological evolution
- 5 Summary

The basic picture



High-mass binary with VHE emission, $\gamma\gamma$ absorption and radio emission.

Estimates: Flux, spectra and morphology

- Flux: $L_{\text{radio}} \sim N_{e^\pm} (E/t_{\text{sync}})_{5 \text{ GHz}} \sim (L_{e^\pm} t_{\text{cool}}/E_{\text{inj}})(E/t_{\text{sync}})_{5 \text{ GHz}}$
where $E_{5 \text{ GHz}}$ comes from $\dot{E}(r)$ ($\sim 10 \text{ mJy}$)
- Spectra: IC cooling ($F_\nu \propto \nu^{-1/2}$), adiabatic cooling (F_ν flat), free free absorption (F_ν harder)
- Size: IC cooling ($X_{\text{radio}} \sim r_{\text{inj}}$), adiabatic cooling
($X_{\text{radio}} \sim r_{\text{inj}} (E_{\text{inj}}/E_{5 \text{ GHz}})^{2/3} \gg r_{\text{inj}}$)

Detailed study

- Monte-Carlo calculations
- Photons are injected in the system and followed till absorbed in the stellar photon field.
- Pairs follow the magnetic field lines (adiabatic invariant approximation) or diffuse anisotropically.
- Pairs evolve suffering IC, synchrotron, adiabatic, relativistic Bremsstrahlung and ionization losses.

Outline


- 1 Introduction
- 2 Exploring the scenario
- 3 Flux and spectral evolution in LS 5039
- 4 Morphological evolution
- 5 Summary

Particular case: LS 5039


- $L_* = 7 \times 10^{38} \text{ erg s}^{-1}$, $T_* = 38000 \text{ K}$, $R_* = 7 \times 10^{11} \text{ cm}$
- $e = 0.33$, $a = 2.2 \times 10^{12} \text{ cm}$
- $i \sim 30^\circ - 60^\circ \rightarrow 45^\circ$
- $v_{w\infty} = 2.5 \times 10^8 \text{ cm s}^{-1}$, $v_{\tan} = 0.1 v_{w\infty}$
- $\dot{M}_w = 3 \times 10^{-7} M_\odot \text{ yr}^{-1}$
- $B_* = 200 \text{ G}$, $B_r = 200 (R_*/r)^{-2} \text{ G}$, $B_\phi = 200 (v_{\tan}/v_{w\infty}) (R_*/r) \text{ G}$
- $D = 0.01 D_{\text{Bohm}}$ ($\delta B(r_g) = 0.1 B$)
- $L_{\gamma > 10 \text{ GeV}} = 10^{36} \text{ erg s}^{-1}$ ($\tau \geq 1$)
- $\Gamma = 2.3$
- Gamma-ray emitter at the compact object location (0th order).

(Casares et al. 2005; Usov & Melrose 1992; Aharonian et al. 2006)

Phase 0.0, ~ SUPC




Spectral energy distribution along the orbit




Outline

- 1 Introduction
- 2 Exploring the scenario
- 3 Flux and spectral evolution in LS 5039
- 4 Morphological evolution
- 5 Summary


Phase 0.0, \sim SUPC; electrons




Phase 0.25




Phase 0.5




Phase 0.75, \sim INFC




Phase 0.0, \sim SUPC




Phase 0.25





Phase 0.5




Phase 0.75, \sim INFC




Imaging the system




Phase 0.0, ~ SUPC; source




Phase 0.25




Phase 0.5



Phase 0.75, ~ INFC




Phase 0.0, ~ SUPC




Phase 0.25


5 GHz Phase 0.25 i: 45 deg [mJy/mas]




Phase 0.5




Phase 0.75, ~ INFC



Phase 0.0, ~ SUPC; image




Phase 0.25



Phase 0.5

5 GHz Phase 0.5 i: 45 deg [mJy/beam]


5
2.6
1.3
0.64
0.32
0.16
0.08




Phase 0.75, ~ INFC

5 GHz Phase 0.5 i: 45 deg [mJy/beam]


5
2.6
1.3
0.64
0.32
0.16
0.08



Phase 0.0, ~ SUPC




Phase 0.25



Phase 0.5

5 GHz Phase 0.5 i: 45 deg [mJy/beam]


5
2.6
1.3
0.64
0.32
0.16
0.08




Phase 0.75, ~ INFC

5 GHz Phase 0.5 i: 45 deg [mJy/beam]


5
2.6
1.3
0.64
0.32
0.16
0.08




Phase 0.0, ~ SUPC; residuals



Phase 0.25




Phase 0.5




Phase 0.75, ~ INFC

5 GHz Phase 0.5 i: 45 deg [mJy/beam]


0.64
0.32
0.16
0.08




Phase 0.0, ~ SUPC




Phase 0.25



Phase 0.5



Phase 0.75, ~ INFC



Outline

1 Introduction

2 Exploring the scenario

3 Flux and spectral evolution in LS 5039

4 Morphological evolution

5 Summary

Summary

- The production of gamma-rays ≥ 100 GeV in massive gamma-ray binaries leads to e^\pm creation.
- Pairs rapidly isotropize and radiate.
- Radio fluxes are detectable.
- Moderate amount of radio variability is expected.
- Radio spectra tend to be flat.
- The structures are resolvable and change significantly along the orbit.
- Important parameters are the magnetic field and the ionization of the wind.
- X-ray-to-radio flux ratios depend strongly on $B(r)$, r_{inj} and Γ .